arXiv:1503.06954v2 [math.NT] 30 Apr 2015

THE ζ-REGULARIZED PRODUCT OVER ALL PRIMES

V.V. SMIRNOV

ABSTRACT. In this paper we prove that the ζ-regularized product over all primes is πe^{μ}, where μ is closely related with the non-trivial zeros of the $\zeta(s)$.

1. INTRODUCTION.

The Chebyshev theta function $\theta(x)$ is defined by

$$\theta(x)=\sum_{p\le x}\log p, \tag{1.1}$$

where p runs over primes. There is a simple relationship between $\theta(x)$ and the prime-counting function $\pi(x)$:

$$\pi(x)=\frac{\theta(x)}{\log x}+\int_2^x\frac{\theta(t)}{t\log^2 t}dt. \tag{1.2}$$

Indeed, let $p_k\in\mathbb{P}$ and let $a_k=\sum_{j=1}^k\log p_j$ so $a_k-a_{k-1}=\log p_k$, hence we have

$$\begin{aligned}\int_2^x\frac{\theta(t)}{t\log^2 t}dt&=\sum_{j=1}^{k-1}\int_{p_j}^{p_{j+1}}\frac{a_j}{t\log^2 t}dt+\int_{p_k}^x\frac{a_k}{t\log^2 t}dt\\&=\sum_{j=1}^{k-1}a_j\left(\frac{1}{\log p_j}-\frac{1}{\log p_{j+1}}\right)+\frac{a_k}{\log p_k}-\frac{a_k}{\log x}\\&=\frac{a_1}{\log p_1}+\sum_{j=2}^{k}\frac{a_j-a_{j-1}}{\log p_j}-\frac{a_k}{\log x}\\&=1+\sum_{j=2}^{k}1-\frac{\theta(x)}{\log x}=\pi(x)-\frac{\theta(x)}{\log x}.\end{aligned}$$

The second Chebyshev function $\psi(x)$ is defined similarly, with the sum extending over all prime powers not exceeding x:

$$\psi(x)=\sum_{p^k\le x}\log p=\sum_{n\le x}\Lambda(n), \tag{1.3}$$

where $\Lambda(n)$ is the von Mangoldt function. $\psi(x)$ can be related to the prime-counting function as follows

$$\Pi(x)=\sum_{n\le x}\Lambda(n)\int_n^x\frac{dt}{t\log^2 t}+\frac{1}{\log x}\sum_{n\le x}\Lambda(n)=\frac{\psi(x)}{\log x}+\int_2^x\frac{\psi(t)}{t\log^2 t}dt. \tag{1.4}$$

$$\Pi(x)=\pi(x)+\frac{1}{2}\pi(x^{1/2})+\frac{1}{3}\pi(x^{1/3})+\cdots. \tag{1.5}$$

Also, $\psi(x)$ has an explicit expression as a sum over the nontrivial zeros of the Riemann zeta function [1, p.104]:

$$\psi_0(x) = \frac{1}{2}\left(\sum_{n \leq x} \Lambda(n) + \sum_{n<x} \Lambda(n)\right) = x - \sum_{\rho} \frac{x^{\rho}}{\rho} - \frac{\zeta'(0)}{\zeta(0)} - \sum_{k=1}^{\infty} \frac{x^{-2k}}{-2k}, \tag{1.6}$$

where ρ runs over the nontrivial zeros of the zeta function. The terms on the right side are arranged by decreasing magnitude. Thus

$$\sum_{p^k \leq x} \log p \sim x \tag{1.7}$$

and the location of the nontrivial zeros of $\zeta(s)$ determines the largest error term. Since the zeros are symmetric about the line $\Re e(s) = \frac{1}{2}$, and since $|x^{\rho}| = x^{\Re e(\rho)}$, the error term is as small as possible if and only if $\Re e(\rho) = \frac{1}{2}$ for all ρ.

2. THE CONSTANT.

The Riemann ξ function is defined as:

$$\xi(s) = \Pi\left(\frac{s}{2}\right)(s-1)\pi^{-s/2}\zeta(s) \tag{2.1}$$

for $s \in \mathbb{C}$, where

$$\Pi(s) = s\Gamma(s) = \Gamma(s+1) = \int_0^{\infty} e^{-t} t^s dt. \tag{2.2}$$

$\xi(s)$ is an entire function – that is, an analytic function of s which is defined for all values of s – and the functional equation of the zeta function is equivalent to $\xi(s) = \xi(1-s)$. Xi-function can also be expanded as infinite product

$$\xi(s) = \xi(0)\prod_{\rho}\left(1 - \frac{s}{\rho}\right), \tag{2.3}$$

where ρ ranges over the roots of the equation $\xi(\rho) = 0$. Since the logarithmic derivative of $\xi(s)$ is on the one hand

$$\sum_{\rho} \frac{d}{ds}\log\left(1 - \frac{s}{\rho}\right) = \sum_{\rho} \frac{1}{s-\rho} \tag{2.4}$$

and on the other hand

$$\frac{d}{ds}\log\Pi\left(\frac{s}{2}\right) - \frac{1}{2}\log\pi + \frac{1}{s-1} + \frac{\zeta'(s)}{\zeta(s)} \tag{2.5}$$

the sum of the series $\sum 1/\rho$ is the value of

$$-\frac{1}{2}\frac{\Pi'(s/2)}{\Pi(s/2)} + \frac{1}{2}\log\pi + \frac{1}{1-s} - \frac{\zeta'(s)}{\zeta(s)} \tag{2.6}$$

at $s = 0$. Now logarithmic differentiation of the product formula for $\Pi(s)$

$$\Pi(s) = \prod_{n=1}^{\infty} \frac{n^{1-s}(n+1)^s}{s+n} = \prod_{n=1}^{\infty}\left(1 + \frac{s}{n}\right)^{-1}\left(1 + \frac{1}{n}\right)^s \tag{2.7}$$

gives

$$\frac{\Pi'(s)}{\Pi(s)} = \sum_{n=1}^{\infty}\left[-\frac{1}{s+n} - \log n + \log(n+1)\right], \tag{2.8}$$

$$-\frac{\Pi'(0)}{\Pi(0)} = \lim_{n\to\infty}\left[1+\frac{1}{2}+\frac{1}{3}+\cdots+\frac{1}{n}-\log(n+1)\right]. \tag{2.9}$$

The number on the right side of this equation is by definition *Euler-Mascheroni constant* and is traditionally denoted γ. Thus [3, §41],[5],[6]

$$\eta := \sum_{\rho}\frac{1}{\rho(1-\rho)} = 2\sum_{\rho}\frac{1}{\rho} = \gamma + 2 - \log 4\pi = 0.04619\ldots \tag{2.10}$$

3. The ζ-regularized product over all primes.

We define the *prime zeta function* $P(s)$, $s=\sigma+i\tau$, through

$$P(s) = \sum_{p} p^{-s} \tag{3.1}$$

with the summation performed over all primes p. The series converges absolutely when $\sigma > 1$. But we also know that [4, p. 70]

$$e^{p^{-s}} = \prod_{n=1}^{+\infty}(1-p^{-ns})^{-\frac{\mu(n)}{n}}, \tag{3.2}$$

where $\mu(n)$ is the Möbius function. So it follows that

$$e^{P(s)} = \prod_{p} e^{p^{-s}} = \prod_{p}\prod_{n=1}^{+\infty}(1-p^{-ns})^{-\frac{\mu(n)}{n}} = \prod_{n=1}^{+\infty}\zeta(ns)^{\frac{\mu(n)}{n}}. \tag{3.3}$$

Taking the logarithmic derivative we get

$$P'(s) = \sum_{n=1}^{+\infty}\frac{\mu(n)}{n}\frac{n\zeta'(ns)}{\zeta(ns)} = \sum_{n=1}^{+\infty}\mu(n)\frac{\zeta'(ns)}{\zeta(ns)}. \tag{3.4}$$

Now consider the integral

$$\begin{aligned}\int_0^\infty \frac{xt^{x-1}}{e^t+1}dt &= \int_0^\infty \frac{xt^{x-1}}{1+e^{-t}}e^{-t}dt \\ &= x\sum_{k=1}^{\infty}(-1)^{k-1}\int_0^\infty t^{x-1}e^{-kt}dt \\ &= x\sum_{k=1}^{\infty}(-1)^{k-1}k^{-x}\int_0^\infty t^{x-1}e^{-t}dt \\ &= (1-2^{1-x})\zeta(x)\Gamma(x+1).\end{aligned} \tag{3.5}$$

Integration by parts gives

$$\lim_{x\to0^+}\int_0^\infty \frac{xt^{x-1}}{e^t+1}dt = \lim_{x\to0^+}\int_0^\infty \frac{t^x e^t}{(e^t+1)^2}dt = \int_1^\infty \frac{du}{(u+1)^2} = \frac{1}{2}. \tag{3.6}$$

Making $x\to 0$ in (3.5) and combining with (3.6), we get $\zeta(0) = -\frac{1}{2}$. We also know that ζ-regularization gives [4, p.71]

$$\sum_{n=1}^{\infty}\mu(n) = -2. \tag{3.7}$$

Therefore

$$P'(0) = \left(\sum_{n=1}^{+\infty} \mu(n)\right) \frac{\zeta'(0)}{\zeta(0)} = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2\log 2\pi. \tag{3.8}$$

Finally, using (2.10), (3.1) and (3.8) we obtain (3.9), where the product is taken over all primes p and $\mu = 2 + \gamma - \eta$.

$$\prod p = \pi e^{\mu} \tag{3.9}$$

References

1. Harold Davenport, Multiplicative Number Theory, *Springer,* (2000).
2. Lowell Schoenfeld, *Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$. II*, Mathematics Of Computation, Volume 30, Number 134, (April 1976), 337–360.
3. Srinivasa Ramanujan, *Highly composite numbers*, Proceedings of the London Mathematical Society, 2, XIV, (1915), 347–409
4. E. Muoz Garca, R. Prez Marco, *The product over all primes is* $4\pi^2$, Commun. Math. Phys. 277, (2008),69–81.
5. Jean-Louis Nicolas, *Petites valeurs de la fonction d'Euler*, J. Number Theory 17, (1983), 375–388.
6. Guy Robin, *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann*, Journal de Mathématiques Pures et Appliquées, Neuvième Série 63 (2), (1984), 187-213.

E-mail address: `ixorg@ya.ru`